

Haldane's Fractional Exclusion Statistics for Multicomponent Systems

Takahiro Fukui and Norio Kawakami

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-01, Japan

(4 August 1994)

Abstract

The idea of fractional exclusion statistics proposed by Haldane is applied to systems with internal degrees of freedom, and its thermodynamics is examined. In case of one dimension, various bulk quantities calculated show that the critical behavior of such systems can be described by $c = 1$ conformal field theories and conformal weights are completely characterized by statistical interactions. It is also found that statistical interactions have intimate relationship with a topological order matrix in Chern-Simons theory for the fractional quantum Hall effect.

05.30.-d, 71.10.+x

It is known that there are possibilities of existing particles which obey fractional statistics in lower dimensions. For example, in the fractional quantum Hall effect (FQHE) quasi-particles carry fractional charges, which may be described by Chern-Simons (CS) theory [1]. In this theory, as well as others, the dimension of space plays a crucial role because fractional statistics is defined on the basis of monodromic properties of wavefunctions. Recently, Haldane [2] has proposed a new definition of fractional statistics independent of the dimension of space. It is based on the state-counting of many-body systems and can be considered as a generalization of the Pauli exclusion principle. Imagine that there is a N -particle system with species $\alpha_i, i = 1, 2, \dots, N$. The N -body wavefunction can be expanded by i th one-particle wavefunction by keeping the coordinates of other particles fixed. Let D_{α_i} be the dimension of such a basis. By observing that D_{α} may be changed when a particle is added without changing the boundary and the size of the system, Haldane has introduced a statistical interaction $g_{\alpha\beta}$ defined by

$$\frac{\partial D_{\alpha}}{\partial N_{\beta}} = -g_{\alpha\beta}, \quad (1)$$

where $g_{\alpha\beta}$ is a constant independent of $\{N_{\alpha}\}$. The special cases of $g_{\alpha\beta} = 0$ and $g_{\alpha\beta} = \delta_{\alpha\beta}$ correspond to free bosons and free fermions, respectively. Since Haldane's work, several authors have investigated it in detail [3–7].

Wu [4] and Bernard and Wu [5] have studied the thermodynamics of single-component particles whose total number of states is given by

$$W = \prod_i \frac{(D_i + N_i - 1)!}{N_i!(D_i - 1)!}, \quad (2)$$

with D_i being a solution of eq.(1), where the species α is regarded as the index i which labels the momentum k_i . They have shown that g_{ij} is given by the two-body S matrix for Bethe ansatz solvable models. As simple examples in one dimension (1D), they calculated g_{ij} for the Bose gas with δ -function interaction and the Calogero-Sutherland model [5]. The former has non-zero off-diagonal elements of g_{ij} which they call mutual statistics, while the latter has only diagonal ones, $g_{ij} = g\delta_{ij}$. Nayak and Wilczek [6] have called such particles

“ g -ons” and examined low-temperature properties in arbitrary dimensions. It may be such g -ons that can be regarded as ideal particles obeying fractional statistics.

In this Letter, we will generalize this approach to systems with internal degrees of freedom. This has been briefly discussed in [5], and a step has been made forward in [7] for the 1D $U \rightarrow \infty$ Hubbard model. First, we will obtain general formulae for thermodynamics, and then calculate bulk quantities in case of 1D to find that critical behavior of such systems can be described by $c = 1$ conformal field theories (CFT). It will next be shown that there is intimate relationship between statistical interactions and a topological order matrix in CS theory for the FQHE [8].

Let us consider a system of N particles whose states are classified by the labels α and i , where $\alpha (= 1, 2, \dots, M)$ and i correspond to internal degrees of freedom and the momentum of particles, respectively. Denoting explicitly these two kinds of indices in eq.(1), the statistical interaction $g_{\alpha i \beta j}$ for the multicomponent system reads,

$$\frac{\partial D_{\alpha i}}{\partial N_{\beta j}} = -g_{\alpha i \beta j}, \quad (3)$$

which results in

$$D_{\alpha i} = - \sum_{\beta j} g_{\alpha i \beta j} N_{\beta j} + D_{\alpha i}^0, \quad (4)$$

where $N_{\alpha i}$ ($D_{\alpha i}$) is the number of particles (holes) labeled by αi , and $D_{\alpha i}^0$ is the number of holes when there are no particles in the systems. Suppose that we have no particles in the system under consideration. Then there are only M kinds of free holes. Therefore, it is natural to choose $D_{\alpha i}^0 = D_{\alpha}^0$ with $D_{\alpha}^0 = D^0$, which we will call a symmetric basis in this paper. Another important choice is $D_{\alpha}^0 = D^0 \delta_{\alpha 1}$, namely, $\alpha = 1$ and $\alpha > 1$ classify charge and internal degrees of freedom, respectively. This is referred to as a hierarchical basis. It is noteworthy that one can find the same bases also in a CS theory for the FQHE [8], as we shall see later. The distribution functions of particles and holes labeled by αi are defined as

$$\rho_{\alpha i} \equiv \frac{N_{\alpha i}}{D^0}, \quad \rho_{\alpha i}^{(h)} \equiv \frac{D_{\alpha i}}{D^0}. \quad (5)$$

Charge for each component is also defined by $t_\alpha \equiv D_\alpha^0/D^0$, which reduces to $\vec{t} = (1, 1, \dots, 1)$ and $(1, 0, \dots, 0)$ in the symmetric and the hierarchical bases, respectively. Note that the quantities (5) play a key role in the fractional exclusion statistics because they are generalizations of Fermi (or Bose) distribution function. In the following, we consider multicomponent systems whose total energy is given by the summation of bare single-particle energies, $E = \sum_{\alpha i} \epsilon_{\alpha i}^0 N_{\alpha i}$, where we set $\epsilon_{\alpha i}^0 \equiv \epsilon_i^0 t_\alpha$. It should be noted that many-body effects are in general incorporated in the distribution of $N_{\alpha i}$.

Let us now examine the thermodynamics of a system whose number of states is given by eq.(2) with replacing $i \rightarrow \alpha i$. Notice that eq.(4) is rewritten as

$$\rho_{\alpha i}^{(h)} + \sum_{\beta j} g_{\alpha i \beta j} \rho_{\beta j} = t_\alpha. \quad (6)$$

It is now straightforward to investigate the thermodynamics following the method of Wu [4] and Bernard and Wu [5], which is based on Yang and Yang's idea [9]. Thermal equilibrium is realized by minimizing the thermodynamic potential $\Omega \equiv E - TS - \sum_\alpha \mu_\alpha N_\alpha$, where $N_\alpha \equiv \sum_i N_{\alpha i}$ and μ_α denote the total number of particles and the chemical potential for the species α , respectively. The entropy S is given by $S = \ln W$ with W replaced $i \rightarrow \alpha i$ in eq.(2). The results are as follows. The free energy F at equilibrium is

$$F = \sum_\alpha \mu_\alpha N_\alpha - T \sum_{\alpha i} D_\alpha^0 \ln[1 + \exp(-\epsilon_{\alpha i}/T)], \quad (7)$$

where the “dressed energy” $\epsilon_{\alpha i} = T \ln(D_{\alpha i}/N_{\alpha i}) = T \ln(\rho_{\alpha i}^{(h)}/\rho_{\alpha i})$ is determined by

$$\epsilon_{\alpha i} - T \sum_{\beta j} \tilde{g}_{\beta j \alpha i} \ln[1 + \exp(-\epsilon_{\beta j}/T)] = \epsilon_{\alpha i}^0 - \mu_\alpha, \quad (8)$$

where $\tilde{g}_{\alpha i \beta j} \equiv g_{\alpha i \beta j} - \delta_{\alpha i \beta j}$. Though we here assume that there is no degeneracy for each state, we can also formulate the thermodynamics by introducing the corresponding weight factor in eqs.(7) and (8) when each state has a certain degeneracy. The formulae (7) and (8) describe thermodynamics of multicomponent systems obeying the fractional exclusion statistics (1).

Now consider low-temperature properties in the thermodynamic limit, which is defined by $N_\alpha \rightarrow \infty$ and $D^0 \rightarrow \infty$ proportionally [5]. Hereafter, we restrict ourselves to 1D systems for simplicity. Then eqs.(6)~(8) read

$$\rho_\alpha^{(t)}(k_\alpha) = t_\alpha - \sum_\beta \int_{-\infty}^{\infty} dk_\beta \tilde{g}_{\alpha\beta}(k_\alpha, k_\beta) \rho_\beta(k_\beta), \quad (9)$$

$$F = \sum_\alpha \mu_\alpha N_\alpha - T \sum_\alpha D_\alpha^0 \int_{-\infty}^{\infty} dk_\alpha \ln[1 + \exp(-\epsilon_\alpha(k_\alpha)/T)], \quad (10)$$

$$\epsilon_\alpha(k_\alpha) = \epsilon_\alpha^0(k_\alpha) - \mu_\alpha + T \sum_\beta \int_{-\infty}^{\infty} dk_\beta \tilde{g}_{\beta\alpha}(k_\beta, k_\alpha) \ln[1 + \exp(-\epsilon_\beta(k_\beta)/T)], \quad (11)$$

where $\rho_\alpha^{(t)} \equiv \rho_\alpha + \rho_\alpha^{(h)}$. The total energy is given by $e \equiv E/D^0 = \sum_\alpha e_\alpha$ with $e_\alpha = \int_{-\infty}^{\infty} dk_\alpha \epsilon_\alpha(k_\alpha) \rho_\alpha(k_\alpha)$. Assume $\tilde{g}_{\alpha\beta}(-k_\alpha, -k_\beta) = \tilde{g}_{\alpha\beta}(k_\alpha, k_\beta)$ and $\epsilon_\alpha^0(-k_\alpha) = \epsilon_\alpha^0(k_\alpha)$. Then at $T = 0$ there exists a “generalized Fermi level” Q_α which satisfies $\epsilon_\alpha(k_\alpha) < 0$ for $|k_\alpha| < Q_\alpha$ and $\epsilon_\alpha(k_\alpha) > 0$ for $|k_\alpha| > Q_\alpha$. Hence $\rho_\alpha(k_\alpha) = 0$ for $|k_\alpha| > Q_\alpha$, and $\rho_\alpha(k_\alpha)$ for $|k_\alpha| < Q_\alpha$ and $\epsilon_\alpha(k_\alpha)$ are given by the solutions of the following equations:

$$\rho_\alpha(k_\alpha) = t_\alpha - \sum_\beta \int_{-Q_\beta}^{Q_\beta} dk_\beta \tilde{g}_{\alpha\beta}(k_\alpha, k_\beta) \rho_\beta(k_\beta), \quad (12)$$

$$\epsilon_\alpha(k_\alpha) = \epsilon_\alpha^0(k_\alpha) - \mu_\alpha - \sum_\beta \int_{-Q_\beta}^{Q_\beta} dk_\beta \epsilon_\beta(k_\beta) \tilde{g}_{\beta\alpha}(k_\beta, k_\alpha). \quad (13)$$

Therefore, in the low-temperature limit, eq.(10) is expanded as

$$F(T) = F(T=0) - \frac{\pi}{6} T^2 (2\pi D^0) \sum_\alpha \frac{1}{v_\alpha} \quad (14)$$

in terms of the Fermi velocity $v_\alpha \equiv \epsilon'_\alpha(Q_\alpha)/\rho_\alpha(Q_\alpha)$. The specific heat in low-temperatures in unit length is then given by $C/(TL) = (\pi/3) \sum_\alpha (1/v_\alpha)$, where $D^0 = L/(2\pi)$ is used [5]. Hence, the critical behavior of 1D systems whose statistical interaction is defined by eq.(3) can be described by M independent $c = 1$ CFT [11]. The responses to “external fields” $\vec{\mu}$ are also calculated as

$$\chi_{\alpha\beta} \equiv \frac{\partial n_\alpha}{\partial \mu_\beta} = 2(\mathcal{Z} V^{-1} \mathcal{Z}^T)_{\alpha\beta}, \quad (15)$$

where $n_\alpha \equiv N_\alpha/D^0 = \int_{-\infty}^{\infty} dk_\alpha \rho_\alpha(k_\alpha)$, $\mathcal{Z}_{\alpha\beta} \equiv Z_{\alpha\beta}(Q_\beta)$ is a boundary-valued “dressed charge” $Z_{\alpha\beta}(k_\beta)$ defined by [10]

$$Z_{\alpha\beta}(k_\beta) = \delta_{\alpha\beta} - \sum_{\gamma} \int_{-Q_\gamma}^{Q_\gamma} dk_\gamma Z_{\alpha\gamma}(k_\gamma) \tilde{g}_{\gamma\beta}(k_\gamma, k_\beta), \quad (16)$$

and a matrix V is defined by $V_{\alpha\beta} \equiv v_\alpha \delta_{\alpha\beta}$. From eq.(15), the compressibility can be obtained as $\chi_c = \sum_{\alpha\beta} t_\alpha \chi_{\alpha\beta} t_\beta$. The excitation spectrum in CFT limit is classified as [10]

$$\Delta e = \frac{1}{4} \vec{\Delta} n (\mathcal{Z}^{-1})^T V \mathcal{Z}^{-1} \vec{\Delta} n + \vec{\Delta} d \mathcal{Z} V \mathcal{Z}^T \vec{\Delta} d \quad (17)$$

except for particle-hole excitations, where $\vec{\Delta} n$ ($\vec{\Delta} d$) is a M -component vector of quantum numbers which change the number of particles (carry the large momentum). This expression also confirms that the critical behavior is described by M independent $c = 1$ gaussian CFT.

Let us now consider a simple system with

$$g_{\alpha i \beta j} = \delta_{ij} G_{\alpha\beta}. \quad (18)$$

The choice (18) implies that the interaction in the momentum space is local, i.e., it takes a δ -function form. We refer to particles obeying (18) as *ideal multicomponent gases with fractional exclusion statistics which is characterized by the statistical interaction $G_{\alpha\beta}$* . One can indeed find that eqs.(6) and (9) reduce to $\rho_\alpha^{(h)}(k) + \sum_\beta G_{\alpha\beta} \rho_\beta(k) = t_\alpha$, which shows an asymmetry between particles and holes as discussed in refs. [12,5,13]. We will later discuss concrete models in 1D which show fractional exclusion statistics (18). From eq.(18), the distribution of particles with species α at finite temperature is given by

$$\rho_\alpha(k) = \sum_\beta G_{\alpha\beta}^{-1}(k, T) t_\beta, \quad (19)$$

where $G_{\alpha\beta}(k, T) \equiv \exp(\epsilon_\alpha(k)/T) \delta_{\alpha\beta} + G_{\alpha\beta}$, and $\epsilon_\alpha(k)$ is a solution of eq.(11), i.e., $\prod_\beta (e^{\epsilon_\beta/T})^{G_{\beta\alpha}} (1 + e^{\epsilon_\beta/T})^{\delta_{\beta\alpha} - G_{\beta\alpha}} = e^{(\epsilon_\beta^0 - \mu_\beta)/T}$. We emphasize that the formula (19) is a generalization of the Fermi (or Bose) distribution function to a multicomponent ideal gas with the statistical interaction $G_{\alpha\beta}$. We here show simple examples in the symmetric basis. First, when $G_{\alpha\beta} = g$, distribution in eq.(19) becomes that of ideal fractional one-component particles, $\rho = \rho^{(g)}$, which has been discussed in [2,3,5,6]. This formula further reduces to $1/(e^{(\epsilon^0 - \mu)/T} - 1)$ for free bosons ($g = 0$) and to $1/(e^{(\epsilon^0 - \mu)/T} + 1)$ for free fermions ($g = 1$). As

another simple example of two-component system with $G_{11} = G_{22} = g$ and $G_{12} = G_{21} = f$, we have $\vec{\rho} = (\rho^{(g')}, \rho^{(g')})$ with $g' \equiv g + f$.

Consider now the $T = 0$ case with general $G_{\alpha\beta}$. There appear M kinds of “generalized Fermi levels” $Q_1, Q_2, \dots, Q_M$. Define the ordering of α by $\tau(\alpha)$, such that $Q_{\tau(1)} \geq Q_{\tau(2)} \geq \dots \geq Q_{\tau(M)}$. Then, in the region $Q_{\tau(n)} > |k| > Q_{\tau(n+1)}$, we find $\rho_{\tau(\alpha)}(k) = \sum_{\beta=1}^n \tau_n(G)_{\alpha\beta}^{-1} t_{\tau(\beta)}$ for $\alpha = 1, \dots, n$ and $\rho_{\tau(\alpha)}(k) = 0$ for $\alpha = n+1, \dots, M$, where $\tau_n(G)$ is a $n \times n$ matrix whose elements are given by $\tau_n(G)_{\alpha\beta} = G_{\tau(\alpha)\tau(\beta)}$ with $\alpha, \beta = 1, 2, \dots, n$. In the absence of external fields one finds, by putting $\mu_\alpha = \mu t_\alpha$, that the ground state is obtained by setting $Q_1 = Q_2 = \dots = Q_M \equiv Q$. The dressed energy $\epsilon_\alpha(k)$, distribution function ρ_α and the density of particles $n_c \equiv N_c/D^0$, where $N_c \equiv \sum_\alpha N_\alpha t_\alpha$, are calculated as

$$\epsilon_\alpha(k) = \sum_\beta G_{\alpha\beta}^{-1} (\epsilon_\beta^0(k) - \mu_\beta), \quad (20)$$

$$\rho_\alpha = \sum_\beta G_{\alpha\beta}^{-1} t_\beta, \quad (21)$$

$$n_c = 2Q\nu, \quad \nu \equiv \sum_{\alpha\beta} t_\alpha G_{\alpha\beta}^{-1} t_\beta. \quad (22)$$

These equations show that Fermi velocities are calculated as $v_\alpha = Q = n_c/(2\nu)$, which are independent of their components. If we take $\epsilon^0(k) = k^2/2$ as the bare spectrum, we then find the ground state energy $e_g = n_c^3/(24\nu^2)$ and the compressibility $\chi_c = 4\nu^2/n_c$. Hence, these bulk quantities are determined solely by ν which does not involve full information of the matrix G . However, if we observe the excitation spectrum $\Delta e = (v/4)\vec{\Delta}nG\vec{\Delta}n + v\vec{\Delta}dG^{-1}\vec{\Delta}d$, there shows up every element of the matrix G , owing to the fact that M kinds of Fermi velocities are the same (See eq.(17)). Here we assume $G^T = G$. Note that this formula for the excitation is of a desirable form required by CFT, and we thus find conformal dimensions determined completely by the matrix G following the idea of finite size-scaling [11].

A remarkable point is that the above matrix G of statistical interactions has close relationship to a topological order matrix in the FQHE. To see this explicitly, it is more convenient to take $\epsilon^0(k) = k$ as the bare spectrum, which only contains right-going particles. It is known that such *chiral* particles in 1D may describe essential properties for edge states of the FQHE [14]. In this case, the excitation spectrum of the present system has

the form $\Delta e = (v/4)\vec{\Delta}nG\vec{\Delta}n$, which may be described by the holomorphic piece of CFT. Recall here that a certain hierarchy of the FQHE can be described by a multicomponent CS Lagrangian $\mathcal{L} = \epsilon^{\mu\nu\lambda}a_{\alpha\mu}K_{\alpha\beta}\partial_\nu a_{\beta\lambda}$ in which the construction of a hierarchy is characterized by the *topological order matrix* K [8]. An important point is that the excitation spectrum obtained here for the chiral model coincides with that for edge states [14] characterized by the topological order matrix K [15]. Therefore we arrive at an instructive consequence that the matrix G of statistical interaction can be identified with a topological order matrix K in CS theory for the FQHE.

For example, the topological-order matrix K used in ref. [8,16] is written in the symmetric basis,

$$K = \sum_{\alpha=1}^M (p_\alpha - 2 + \delta_{\alpha 1}) C_M^{(\alpha)} + I \quad (23)$$

with an odd integer p_1 and even integers p_α ($\alpha > 1$), where I is a unit matrix and $C_M^{(\alpha)}$ is a $M \times M$ matrix defined by $(C_M^{(\alpha)})_{ij} = 1$ for $i, j = \alpha, \alpha + 1, \dots, M$ and $= 0$ for otherwise. Substituting eq.(23) into eq.(22), we have

$$\nu = \frac{1}{p_1 - \frac{1}{p_2 - \dots \frac{1}{p_M}}}. \quad (24)$$

Note that this quantity is nothing but the filling factor for the FQHE described by (23). It is known [8] that the symmetric basis is converted into the hierarchical basis by the matrix W defined by $W_{\alpha\beta} = \delta_{\alpha\beta} - \delta_{\alpha+1,\beta}$ as follows; $\vec{t} \rightarrow \vec{t}W$ and $K \rightarrow W^T K W$. In this basis, distribution functions (21) are easily calculated at zero temperature; $\rho_\alpha = \prod_{\beta=1}^\alpha \nu_\beta$, where $\nu_\beta = \sum_{i=1}^\beta G_{i1}^{(\beta)-1}$ with $\beta \times \beta$ matrix $G^{(\beta)}$ defined by $G_{ij}^{(\beta)} \equiv G_{M-\beta+i, M-\beta+j}$. In particular, when $p_1 = 2m + 1$ and $p_\alpha = 2$ for $\alpha = 2, 3, \dots, M$, we have the filling factor $\nu = M/(2mM + 1)$ and the corresponding distribution functions $\vec{\rho} = (\nu, (M-1)/M, (M-2)/(M-1), \dots, 2/3, 1/2)$. This family corresponds to a fundamental hierarchy which has been observed in the FQHE [1].

Finally some comments are in order for concrete models in 1D, which exhibit essential

properties of fractional exclusion statistics. It has been claimed that *ideal* particles with fractional exclusion statistics are realized by 1D quantum models with $1/r^2$ interaction [17,18] and their multicomponent versions [19]. In fact, this fractional property has been shown explicitly for the Haldane-Shastry spin chain [20] and also for the Calogero-Sutherland model [12,13,21]. For cases with internal degrees of freedom discussed here, we find that eqs.(6)~(8) indeed reduce to the Bethe ansatz equations for multicomponent Calogero-Sutherland model with a special coupling [22], if we convert the expressions to those of the hierarchical basis. Hence, an ideal multicomponent gas with statistical interaction G may also be realized by multicomponent $1/r^2$ models.

On the other hand, for interacting 1D particles with general type of interaction, excitations cannot be regarded as ideal particles. Nevertheless, if we restrict ourselves to the CFT (low-energy) limit of such systems, the excitation can be well approximated by ideal particles with fractional exclusion statistics. For example, consider the 1D Hubbard model in the CFT limit. There are two elementary modes called spinon and holon, which have different velocities $v_s \neq v_h$ in general. Hence, it seems difficult to deduce the matrix G from eq.(17). However, these velocities are not universal, and therefore by choosing $v_s = v_h$ hypothetically, we can conclude that statistical interaction G is given by $G = I + \alpha C_2^{(1)}$ with $\alpha \equiv (1/K_\rho - 1)/2$, where K_ρ ($1/2 \leq K_\rho \leq 1$) is the critical exponent (divided by 4) for the $4k_F$ oscillating part of the density-density correlation function [23]. This kind of interpretation for the elementary excitation is valid in general for the class of multicomponent Luttinger liquids.

In summary, we have applied the idea of Haldane's fractional statistics to systems with internal degrees of freedom. The critical behavior of such 1D systems can be described by multiple $c = 1$ CFT with conformal weights given by statistical interaction G . It has also been shown that there is intimate relationship between statistical interactions and a topological order matrix in CS theory for the FQHE. It is quite interesting and still open to study how the fractional exclusion statistics control the physics of correlated particles in higher dimensions.

This work is supported by the Grant-in-Aid from the Ministry of Education, Science and Culture.

REFERENCES

- [1] *The Quantum Hall Effect*, eds. R.E. Prange and S.M. Girvin, Springer-Verlag 1990:
Quantum Hall Effect, ed. M. Stone, World Scientific, 1992.
- [2] F. D. M. Haldane, Phys. Rev. Lett. **67**, 937 (1991).
- [3] M.V.N. Murthy and R. Shankar, Phys. Rev. Lett. **72**, 3629 (1994); preprint IMSc-94/24,
cond-mat/9404096.
- [4] Y.-S. Wu, preprint UU-HEP/94-1, cond-mat/9402008.
- [5] D. Bernard and Y.-S. Wu, preprint SPhT-94-043, UU-HEP/94-03, cond-mat/9404025.
- [6] C. Nayak and F. Wilczek, preprint PUPT 1466, IASSNS 94/25, cond-mat/9405017.
- [7] K. Byczuk and J. Spalek, preprint cond-mat/9406095.
- [8] X.G. Wen and A. Zee, Phys. Rev. **B46**, 2299 (1992).
- [9] C.N. Yang and C.P. Yang, J. Math. Phys. **10**, 1115 (1969).
- [10] A.G. Izergin, V.E. Korepin and N.Y. Reshetikhin, J. Phys. **A22**, 2615 (1989).
- [11] H.W.J. Blöte, J.L. Cardy and M.P. Nightingale, Phys. Rev. Lett. **56**, 742 (1986): I.
Affleck, Phys. Rev. Lett. **56**, 746 (1986).
- [12] B. Sutherland, in *Lecture Notes in Physics* **242**, Springer-Verlag, 1985.
- [13] Z.N.C. Ha, preprint IASSNS-HEP-94/27, cond-mat/9405063.
- [14] X.G. Wen, Phys. Rev. **B41**, 12835 (1990); J. Mod. Phys. **B6**, 1711 (1992): M. Stone,
Ann. Phys. **207**, 38 (1991).
- [15] Except for velocities which are non-universal quantities depending on a shape of FQHE
droplet.
- [16] B. Blok and X.G. Wen, Phys. Rev. **B42**, 8133 (1990): N. Read, Phys. Rev. Lett. **65**,

- 1502 (1990).
- [17] F. Calogero, J. Math. Phys. **10**, 2197 (1969): B. Sutherland, J. Math. Phys. **12**, 246, 256 (1971); Phys. Rev. **A4**, 2019 (1971); **A5**, 1372 (1971).
 - [18] F.D.M. Haldane, Phys. Rev. Lett. **60**, 635 (1988): B.S. Shastry, Phys. Rev. Lett. **60**, 639 (1988).
 - [19] Y. Kuramoto and H. Yokoyama, Phys. Rev. Lett. **67**, 1338 (1991): N. Kawakami, Phys. Rev. **46**, 1005 (1992): Z.N.C. Ha and F.D.M. Haldane, Phys. Rev. **B46**, 9359 (1992): K. Hikami and M. Wadati, J. Phys. Soc. Jpn, **62**, 469 (1993): J.A. Minahan and A.P. Polychronakos, Phys. Lett. **B302**, 265 (1993).
 - [20] F.D.M. Haldane, Z.N.C. Ha, J.C. Talstra, D. Bernard and V. Pasquier, Phys. Rev. Lett. **69**, 2021 (1992): F.D.M. Haldane, in *Proc. of the 16th. Taniguchi Symposium on Condensed Matter Physics*, eds. A. Okiji and N. Kawakami, Springer-Verlag, 1994: Z.N.C. Ha and F.D.M. Haldane, preprint IASSNS-HEP-94/42, cond-mat/9406059.
 - [21] V. Pasquier, preprint SPhT/94-060, hep-th/9405104.
 - [22] N. Kawakami, Phys. Rev. Lett. **71**, 275 (1993); Prog. Theor. Phys. **91**, 189 (1994).
 - [23] H.J. Schulz, Phys. Rev. Lett. **64**, 2831 (1990): N. Kawakami and S.-K. Yang, Phys. Lett. **148A**, 359 (1990). H. Frahm and V.E. Korepin, Phys. Rev, **B42**, 10553 (1990):